



**Measure Abbreviation:** CARD 02 (QCDR Measure ID: ASPIRE18)

**Data Collection Method:** This measure is calculated based on data extracted from the electronic medical record combined with administrative data sources such as professional fee and discharge diagnoses data. This measure is explicitly not based on provider self-attestation.

**Description:** Percentage of cases without elevated postoperative Troponin I levels (>0.60) or documentation of perioperative myocardial injury.

**NQS Domain:** Effective Clinical Care

**Measure Type:** Outcome

**Scope:** Calculated on a per case basis.

**Measure Summary:** CARD02 is an outcome measure that identifies patients that had elevated troponin levels (Troponin I > 0.6) within 72 hours postoperatively. Troponin I levels are accurate markers of myocardial infarction.

**Rationale:** Postoperative myocardial infarction within 72 hours (as defined by a Troponin I level >3.6 times the 99<sup>th</sup> percentile upper reference limit, usually no greater than 1.00 ng/mL)<sup>1,2</sup> is associated with a significantly increased risk of 30-day mortality. Furthermore, any amount of postoperative myocardial injury (as defined by a Troponin I level >0.03 ng/mL) is an independent predictor of 30-day mortality.<sup>3</sup> Preventing myocardial infarction is an important anesthetic goal.

**Inclusions:** All anesthetic cases.

**Exclusions:**

- ASA 5 and 6 cases.
- Outpatient cases.
- Troponin I  $\geq$  0.01 within 42 days prior to anesthesia start.\*
- Pacemaker insertions (CPT: 00530)
- Cardiac Ablation (CPT: 00537)
- Cardiac surgery without pump (CPT: 00560)
- Cardiac surgery with pump and <1 year old (CPT: 00561)
- Cardiac surgery with pump and > 1 year old (CPT: 00562)
- Cardiac surgery with hypothermic arrest (CPT: 00563)
- CABG without pump (CPT: 00566)
- CABG with pump (CPT: 00567)
- Heart Transplant (CPT: 00580)
- Testing of cardioversion or defibrillator functions (CPT: 00534)
- Cardiac catheterization (CPT: 01920)

## (CARD02) Measure Specification Page 2 of 5

\*Rationale for excluding patient with troponin elevation within 42 days prior to date of surgery is based upon ACC/AHA guidelines recommending a delay in elective surgery for 6 weeks following myocardial infarction.<sup>4</sup>

### MPOG Concept IDs Required:

Myocardial Injury MPOG Concept IDs		Troponin MPOG Concept ID	
90201	CPOM measure Cardiac Arrest	5011	Formal lab- Troponin
90202	CPOM measure Myocardial Ischemia		

### Data Diagnostics Affected:

- Percentage of Cases with Postoperative Troponin (Postoperative Troponin)
- Percentage of Cases with Professional Fee Procedure Codes (Pro Fee Procedures)
- Percentage of CPT Codes from Anesthesia Professional Fee Billing that are actually Anesthesia Codes (Anesthesia Codes)
- Percentage of Cases with a Meaningful Admission Type Mapping (Admission Type Mapping)
- Percentage of Cases with ASA Status (Cases with ASA Status)

### Collations Used:

- Anesthesia Start
- Anesthesia End
- ASA 5 or 6
- BP 01

### Other Measure Build Details:

- If another case starts within 72 hours, then the time window ends at anesthesia start of the subsequent case.
- For the preoperative Troponin I exclusion, cases with preoperative Troponin I values with 'less than (<)' included in the result will be included up to '<0.31.' For example, preoperative Troponin I levels that are resulted as '<0.02' will be included for the measure. However, a preoperative Troponin I value of 0.02 will be excluded. The rationale for this is each pathology department determines the lower bound for detecting Troponin I levels accurately. This is standardized to the health system but is not standard across all participating sites.
- Algorithm for determining Case Duration:  
Case Start:
  1. Anesthesia Induction End. If not available, then
  2. Anesthesia Induction Begin. If not available, then
  3. Procedure Start. If not available, then
  4. Patient in Room. If not available, then
  5. Anesthesia Start

**Case End:**

1. Patient Extubated. If not available, then
2. Procedure End. If not available, then
3. Patient Out of Room. If not available, then
4. Anesthesia End.

**Success:**

In cases with a Troponin I value(s) available within 72 hours after anesthesia end, all values must be less than or equal to 0.6 for the case to pass.

If no Troponin I values are available within 72 hours after anesthesia end and there is no documentation of perioperative myocardial injury (MPOG Concepts: 90201, 90202), the case will pass.

**Threshold:** 95%.

**Responsible Provider:** Providers assigned to patient longest duration of case unless there are providers who failed BP 01 during case. In that case, BP 01 failure takes precedence over longest duration. See 'Other Measure Build Details' section of this specification to view the algorithm used for determining case duration.

**Method for determining Responsible Provider:**

- 1) Provider(s) who failed BP 01. If not applicable,
- 2) Provider(s) signed into the case for the longest duration.

**Risk Adjustment (for outcome measures):**

*To evaluate provider-level risk adjustment we will calculate the observed to expected outcomes ratio (O/E). The O/E is calculated using a logistic regression model and predicts (given a set list of dependent patient and hospital level variables) the expected probability of having an elevated Troponin I level. We adjust for surgery risk score, emergent procedures, ASA, gender, age, body mass index, laboratory values, and teaching versus private hospital. Patient specific comorbidities are evaluated as well.*

**References:**

1. Thygesen K, Alpert JS, Jaffe AS, Simoons ML, Chaitman BR, White HD. Third universal definition of myocardial infarction. *Global Heart*. 2012;7(4):275-295.
2. Devereaux PJ, Xavier D, Pogue J, et al. Characteristics and short-term prognosis of perioperative myocardial infarction in patients undergoing noncardiac surgery: a cohort study. *Annals of internal medicine*. 2011;154(8):523-528.
3. Botto F, Alonso-Coello P, Chan MT, et al. Myocardial injury after noncardiac surgery: a large, international, prospective cohort study establishing diagnostic criteria, characteristics, predictors, and 30-day outcomes. *Anesthesiology*. 2014;120(3):564-578.
4. Fleisher LA, Fleischmann KE, Auerbach AD, et al. 2014 ACC/AHA guideline on perioperative cardiovascular evaluation and management of patients undergoing noncardiac surgery. *Journal*

*of nuclear cardiology: official publication of the American Society of Nuclear Cardiology.*  
2015;22(1):162-215.

**CARD 02 Flow Diagram**

